

NG Agriculture Insights

A Popular Magazine (ISSN: 3107-5703)
Volume 2, Issue 5, September- October, 2026, 134- 137

Canopy Architecture Management in Fruit Crops

Neeharika Kanth[1*], Jyotsnarani Pradhan[2] & Kundan Kishore[3]

[1]Assistant Professor (Fruit Science), Pandit Deen Dayal Upadhyay College of Horticulture and Forestry, RPCAU, Piprakothi, Bihar-845429, India

[2]Assistant Professor (Botany, Plant Physiology and Biochemistry), College of Basic Sciences & Humanities, RPCAU, Pusa, Samastpur, Bihar- 848125, India

[3]Dean, Pandit Deen Dayal Upadhyay College of Horticulture and Forestry, RPCAU, Piprakothi, Bihar-845429, India

*Corresponding author email ID: neeharika@rpcau.ac.in

A B S T R A C T

Canopy architecture management is an important component of modern fruit production, influencing light interception, photosynthetic efficiency, vegetative growth, flowering, fruit set, fruit quality and orchard productivity. The canopy architecture management includes spatial arrangement of trunk, scaffold branches, shoots and leaves which determines the effective distribution of solar radiation inside plant canopy. The poorly or unmanaged canopies lead to reduction in overall production of crops. This happens due to insufficient penetration of light inside crop canopy, hence, causing hindrance in sprays and increased risk of disease, insects and pests. Appropriate training, pruning, branch bending, crop-load regulation, rootstock selection and high-density planting can help maintain a balanced canopy. The selection of canopy manipulation strategy depends on growth habit of trees, fruiting behavior, cultivar, rootstock and agroclimatic conditions.

Keywords: *Canopy Architecture, Training, Pruning, Light Interception, High-Density Planting, Fruit Quality, Orchard Management*

Submitted: 15-10-2026 Accepted: 22-10-2026 Online: 29-10-2026

1. INTRODUCTION

Fruit trees, being perennial in nature needs canopy architecture management as it is a long term venture. The spatial distribution of shoots, branches and leaves is an important aspect of canopy architecture management. Canopy architecture management involves modifying and maintaining this structure to improve the efficiency of light interception, facilitate orchard operations and sustain fruit production over the life of the orchard. Modern orchard systems emphasize on planned tree architecture, suitable plant density, regular training and pruning from early phase of crop establishment (Singh, 2018). The relationship between canopy structure and productivity is especially important in mango, guava, citrus, litchi, apple, peach and other perennial fruit crops. But a single canopy management strategy doesn't hold good for every crop.

https://doi.org/10.5281/zenodo.23753823

© 2026 The Author(s). This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0).

2. PHYSIOLOGY OF CANOPY ARCHITECTURE MANAGEMENT

Plant physiology of canopy architectural management is all about the link between plant anatomy, light absorption, photosynthesis, and productivity. The well-managed canopy will ensure efficient distribution of light, airflow, and nutrients in the plants. It results in effective flowering, fruiting, development of fruits, and increased yields. This knowledge of the physiology of the processes enables one to design proper methods of canopy management.

2.1 Role of light interception

Light is the primary energy source for photosynthesis, through which leaves produce carbohydrates required for shoot growth, root development, flowering and fruit development. The spatial distribution of light inside canopy plays major role in amount of photosynthesis, incidence of attack of diseases and insects and overall growth of plants. The dense canopies restrict proper light penetration inside canopy and affect overall plant growth and effect fruit production as well as quality. Thus, canopy management aims to achieve a favorable balance between total and uniform light interception in canopy (Zaffer *et al*., 2020; Breen *et al*., 2021).

2.2 Vegetative and reproductive growth balance

The excessive vegetative growth hinders the accumulated photosynthates to divert to the reproductive sink of the trees. The proper training and pruning of unproductive or intermingling branches make balance between vegetative and reproductive growth of trees. The response to pruning depends on its severity, timing, crop species and physiological condition of the tree (Patel *et al*., 2024; Dhurve *et al*., 2025).

2.3 Canopy microclimate and crop health

Canopy density has major effect on temperature, humidity, air movement inside canopy. The increased humidity and temperature inside unmanaged canopy attract insects, pests and diseases which is detrimental for tree growth and health. Appropriate canopy architecture can also improve light exposure, fruit color development, harvesting accessibility and the effectiveness of irrigation, fertilization and plant-protection operations (Patel *et al*., 2024).

3. MAJOR TECHNIQUES FOR CANOPY ARCHITECTURE MANAGEMENT

3.1 Training and pruning of fruit trees

Training is the process of developing a desired tree framework by directing the growth and positioning of the trunk, scaffold branches and shoots. It is particularly important during the establishment phase because the initial framework influences canopy development and orchard management for many subsequent years.

a) Open- center or vase system

The central leader is removed or suppressed to encourage three to five well-spaced scaffold branches. It results in an open canopy that promotes penetration of sunlight and airflow. This system is widely adopted in peaches and other fruit trees and can be modified for particular varieties of guava and other fruit plants.

b) Central-leader system

An adequate central stem is retained along with proper lateral branches spacing. This tree form is appropriate for an apple as well as some other fruits, which require a vertical growth form of canopy. Tree vigor and height should be controlled depending on the rootstock and plant density.

c. Modified leader

An effective framework is established by controlling excess vertical growth and overcrowding. This technique can be applied to several tropical and subtropical species of trees such as mangoes where structural integrity of the canopy should be maintained without cutting off productive branches.

d. Espalier and trellis systems

Branches are trained along wires or a support framework to develop a relatively planar canopy. These systems can improve canopy accessibility, light distribution and operational efficiency in suitable high-density orchards. Their success depends on appropriate cultivars, rootstocks and continued training (Mir *et al*., 2020)

Pruning involves the selective removal of shoots, branches or other plant parts to maintain the desired canopy structure. It is one of the most widely used methods for managing tree vigor, improving light distribution and maintaining productive fruiting wood. The principal pruning methods include thinning of entire shoots which are intermingling or making canopy over crowded. Heading back includes topping of vertical growth of trees in order to break apical dominance and cause lateral growth and flowering. Removal of dead and diseased wood: Damaged, dead, crossing and infected branches are removed to improve tree health and facilitate orchard

operations. In summer pruning, selected shoots are removed during active growth to regulate canopy density and manage unwanted vegetative growth. In case of rejuvenation pruning selected old, unproductive or overcrowded branches are removed or shortened to rebuild the canopy and encourage productive new growth (Kumar *et al*., 2015).

3.2 Branch bending

This technique involves repositioning of upright shoots to a more horizontal inclined position. Branch bending is an important canopy-management technique in which upright shoots are repositioned to a more horizontal or moderately inclined orientation. It can reduce apical dominance, encourage the development of lateral shoots and influence the formation of reproductive structures in certain crops (Dhurve et al., 2025).

3.3 High-density planting and canopy size regulation

High-density planting involves establishing more trees per unit area and maintaining their size and shape through suitable rootstocks, spacing, training and pruning. Its purpose is not simply to increase plant population but to optimize the distribution of productive foliage within the available orchard space.

3.4 Use of rootstock and plant growth regulators

The rootstock also affects the vigour, canopy size, precociousness and adaptation of the tree to soil and climate. Size controlling rootstocks allow easier maintenance of smaller canopies and planting of high density plantings, especially in fruit crops like apple.

PGRs may also be applied in certain fruit crops for the control of growth or physiological phenomena such as flowering. Paclobutrazol has yielded encouraging results in regulating growth and flowering in mango. (Dhurve *et al*., 2025).

3.5 Role of canopy management in fruit yield and quality

Canopy architecture influences fruit size, color development, soluble solids and the uniformity of maturation through its effects on light exposure and source–sink relationships. A well-managed canopy can improve the proportion of foliage and fruit-bearing sites receiving adequate light. It can also reduce competition between adjacent trees and facilitate more uniform fruit development (Breen et al., 2021). Manageable tree height and well-spaced branches facilitate pruning, spraying, fruit thinning and harvesting. They may also improve the reach and effectiveness of plant-protection equipment and reduce the need for hazardous climbing during harvest (Singh, 2018). Old fruit trees may become unproductive because of excessive height, overcrowding, declining productive wood or prolonged neglect. Rejuvenation through selective or staged pruning, canopy rebuilding and, where appropriate, top-working can restore a more functional tree structure.

4. EMERGING TECHNOLOGIES IN CANOPY ARCHITECTURE MANAGEMENT

New technologies that help manage canopy architecture are revolutionizing the process of monitoring and regulating plant development, canopy structure, and crop production. Technologies like remote sensing, drones, artificial intelligence, and precise pruning methods allow for efficient management of the canopy depending on the needs of plants at any moment. These technologies ensure resource efficiency and high-quality fruits.

4.1. Canopy mapping in three dimensions

Digital images and LIDAR or photogrammetry have the potential to describe tree height, canopy volume, foliage pattern and gaps. Such information could be helpful for detecting overcrowded areas and the impact of pruning operations.

4.2 Remote sensing and artificial intelligence

Imagery from either ground-based cameras or UAVs could potentially aid in determining canopy vigour, locating gaps, measuring crop load and detecting variations in orchard development. The practicality of such imagery will depend on its quality, crop architecture, algorithms and field calibration.

4.3 Precision pruning

Digital canopy maps and decision support systems could help identify branches that need pruning and determine how much pruning is needed. Automated recommendations need to take into consideration the fruiting characteristics of the tree, its health and future training system and not only canopy density.

4.4 Precision irrigation and nutrition management

Information about canopy size and vigour, when combined with soil and weather measurements as well as plant condition information, can help make decisions about irrigation and fertilization needs.

5. CONCLUSION

Canopy management is one of the most essential techniques for optimizing productivity, quality, and efficiency in fruit orchards. Through managing the structure of the trees by utilizing training, pruning, branching, choosing rootstocks, and planting density, growers will be able to manage light interception, retain productive fruiting wood, and manage orchard operations. Canopy management in the future should incorporate current practices, knowledge, and technology specific to the crop.

REFERENCES

Breen, K. C., Tustin, D. S., van Hooijdonk, B. M., Stanley, C. J., Scofield, C., Wilson, J. M., Oliver, M. J., & Dayatilake, G. A. (2021). Use of physiological principles to guide precision orchard management and facilitate increased yields of premium quality fruit. *Acta Horticulturae*, 1314. https://doi.org/10.17660/ActaHortic.2021.1314.31

Dhurve, L., Kumar, A., & Joseph, A. V. (2025). Advances in canopy management for enhancing productivity and sustainability in perennial fruit crops: A comprehensive review. *International Journal of Environment and Climate Change*, 15(11), 242–250. https://doi.org/10.9734/ijecc/2025/v15i115110

Kumar, R., Kumar, A., Kumar, V., & colleagues. (2015). Rejuvenating old, senile orchards of tropical and subtropical fruits for enhanced production and improved quality: A review. *The Indian Journal of Agricultural Sciences*, 85(3). https://doi.org/10.56093/ijas.v85i3.47057

Mir, J. I., Ahmed, N., Singh, D. B., Sharma, O. C., Raja, W. H., Shafi, W., Zaffer, S., Jan, S., & Kirmani, S. N. (2020). Canopy management to improve fruit quality of Coe Red Fuji, Granny Smith and Spartan varieties of apple (Malus domestica). *The Indian Journal of Agricultural Sciences*, 90(8). https://doi.org/10.56093/ijas.v90i8.105902

Patel, S. K., Kumar, V., Mishra, D., Kumar, A., Kumar, P., Dansena, D., Ali, I., Patel, B., & Sonkar, A. (2024). Tree canopy architecture and engineering: Physiological and molecular aspects. *Journal of Scientific Research and Reports*, 30(4), 89–94. https://doi.org/10.9734/jsrr/2024/v30i41892

Singh, V. K. (2018). High-density planting system in subtropical fruit crops. *Indian Horticulture*, 63(1). https://epubs.icar.org.in/index.php/IndHort/article/view/84612